

A search for magnetic stars in late stages of stellar evolution

V.G. Elkin

*Special Astrophysical Observatory, Nizhnij Arkhyz, Karachaevo-Cherkesia,
Russia*

Two well-known types of stars with strong dipolar magnetic fields have been extensively studied. These are the magnetic CP stars and the magnetic white dwarfs. In a number of papers, it is suggested that these types of stars may be related by evolution: CP stars could be progenitors of magnetic white dwarfs (Angel et al. 1981). Investigation showed that magnetic fields in CP stars do not change significantly during their lifetime on the main sequence (Glagolevskij et al. 1986).

What happened with the magnetic field later on, after the star has left the main sequence? Stępień (1993) proposed that some very slowly rotating yellow giants or subgiants with strong chromospheric and spot activity, could be former Ap stars with strong magnetic fields.

When the stars become red giants, they undergo a deep convective mixing (first dredge-up) at the red giant branch (RGB), before helium ignition in the core. At the asymptotic giant branch (AGB), they have a second dredge-up (Iben 1991). The internal structure of the star significantly changes in this case.

After the RGB and AGB, the star loses a large part of its own mass. If the field is not completely destroyed at these episodes, it must also change much. This can be traced if the magnetic fields are investigated in stars of intermediate evolution stages, between the main sequence and white dwarfs.

In this context, it is our opinion that there are two types of stars which should be examined for the presence of strong dipolar magnetic fields: the horizontal branch A and B stars, and the hot subdwarfs. These stars are appropriate for direct magnetic measurements using Zeeman spectroscopy, since they show strong and narrow absorption lines in their spectra. In addition, some of them may possess chemical abundance anomalies similar to main sequence CP stars (Bashek and Sargent 1976, Kodaira and Philip 1984, Heber 1992), suggesting an intriguing link between the CP stars and the evolved A and B stars.

Here, we present the results of a continuing search for strong magnetic fields in horizontal branch stars and in hot subdwarfs, using Zeeman spectra obtained at the 6-metre telescope.

References

Angel, J.R.P., Landstreet, J.D., and Borra E.F.: 1981, *Astrophys. J., Suppl. Ser.* **45**, 457

Contrib. Astron. Obs. Skalnaté Pleso ?, (2008), 1– 3.

Table 1. Stellar parameters of Field Horizontal Branch stars

Star	M_v	T_{eff}	$\log g$	$v \sin i$	[Fe/H]	$M/M_{\odot}$
HD 60778	0.95	8200	3.2	17	-0.5	0.48
HD 74721	1.0	8700	3.6	9	-1.11	
HD 86986	0.9	7900	3.1	20	-1.3	
HD 97859		15300	4.0	105	-1.2	
HD 109995	1.1	8300	3.2	15	-1.3	0.37
HD 117880	2.5	8400	3.6		-1.7	0.42
HD 161817	0.8	7500	2.9	10	-1.0	
HD 169027	0.3	11600	3.8			

Table 2. Magnetic fields and radial velocities of FHB stars

Star	B_e (G)	$\pm \sigma$ (G)	RV (km/s)
HD 60778	+150	115	+74
HD 74721	+240	150	+40
HD 86986	-430	580	+3
HD 97859	-400		+66
HD 109995	-820	470	
	+700	520	-139
HD 117880			+142
HD 161817	+30	110	
	-550	140	-368
	-90	80	
	+100	160	-366
HD 169027	-320	820	-26

- Bashek, B., and Sargent A.I.: 1976, *Astron. Astrophys.* **53**, 47
- Glagolevskij, Yu.V., Klochkova V.G., Kopylov I.M.: 1986, in *Upper Main Sequence Stars with the Anomalous Abundances*, ed.: C. R. Cowley, M. M. Dworetsky, and C. Megessier, , Dordrecht: Reidel, 29
- Heber, U.: 1992, in *The Atmospheres of Early-Type Stars*, ed.: U. Heber, C.S. Jeffery, Lecture Notes in Physics **401**, Springer: Berlin, 233
- Iben, I.: 1991, *Astrophys. J., Suppl. Ser.* **76**, 55
- Kodaira, K., and Philip, A.G.D.: 1984, *Astrophys. J.* **278**, 202
- Śtepień: 1993, *Astrophys. J.* **416**, 368

Table 3. Parameters for hot subdwarfs

Star	Sp	M_v	T_{eff}	log g
HD 4539	sdB	3.6	25000	5.4
Feige 87	sdB		28000	5.6
HD 76431	sdB	2.0	35000	4.5
BD+75°325	sdO	4.0	50000	5.3
BD+25°2534	sdOp	4.1	34000	5.5
BD+25°4655	sdO		42200	6.7
HD 128220	sdO+G		42500	4.5
HD 149382	sdOB	4.5	35000	5.5

Table 4. Magnetic field measurements of hot subdwarfs

Star	B_e (G)	$\pm\sigma$ (G)	Comment
HD 4539	-1300 +670	2100 500	$\lambda 4000 - 4600 \text{ \AA}$ <i>HeI</i> $\lambda 5876$ 2 spectra
Feige 87	-1800	3500	H-magnetometer.
HD 76431	-50	130	$\lambda 4000 - 4600 \text{ \AA}$
BD + 75°325	+1260 -1680 +970 -80	870 60 140 280	$\lambda 4000 - 4600 \text{ \AA}$ <i>HeI</i> $\lambda 5876$ 3 spectra <i>HeI</i> $\lambda 4471$ 3 spectra <i>HeI</i> $\lambda 5876$ 2 spectra
BD + 25°2534	+460 +1750 -1300 -1100 -100	 230 600 390 400	<i>H</i> α 1 spectrum <i>H</i> α + <i>HeI</i> $\lambda 5876$ <i>HeI</i> $\lambda 5876$ 2 spectra <i>HeI</i> $\lambda 5876$ 2 spectra <i>HeI</i> $\lambda 5876, 6678$
BD+25°4655	+240 400 -400	340 280 240	<i>HeI</i> $\lambda 6678$ 2 spectra $\lambda 4400 - 4500 \text{ \AA}$ $\lambda 4400 - 4500 \text{ \AA}$
HD 128220	-520 -340 170	950 400	$\lambda 4000 - 4600 \text{ \AA}$ <i>HeI</i> $\lambda 5876$ 3 spectra <i>HeI</i> $\lambda 5876$ 1 spectrum
HD 149382	-10 -1200	890 900	$\lambda 4000 - 4600 \text{ \AA}$ <i>HeI</i> $\lambda 5876$ 3 spectra